



Professor of Orthopaedic Surgery Division of Sports Medicine Tel: (646) 501-7223

Rehabilitation Protocol: Arthroscopic Meniscus Repair

name: Date:
Diagnosis: Date of Surgery:
Phase I (Weeks 0-6)
Weight bearing: Partial weight bearing with crutches weeks 0-4, WBAT after week 4
Hinged Knee Brace: worn for 6 weeks post-op
 Locked in full extension for ambulation and sleeping – remove for hygiene and PT (Weeks 0-1) Unlocked for ambulation and removed while sleeping, for hygiene and PT (Weeks 1-4)
 Range of Motion – AAROM → AROM as tolerated
 Weeks 0-4: Full ROM – No weight bearing at flexion angles greater than 90° Weeks 4-6: Full ROM as tolerated – progress to flexion angles greater than 90°
Therapeutic Exercises
 Quad/Hamstring sets, heel slides, straight leg raises, co-contractions Isometric abduction and adduction exercises
O Patellar Mobilizations
O At 4 Weeks : can begin partial wall-sits – keep knee flexion angle less than 90°
Phase II (Weeks 6-12)
• Weight bearing: As tolerated discontinue crutch use at 6 weeks
Hinged Knee Brace: Discontinue brace use when patient has achieved full extension with no evidence of the continue brace.
extension lag
Range of Motion – Full active ROM
Therapeutic Exercises
Closed chain extension exercises, Hamstring strengthening Lyngag 0.00% Lagranges 0.00%
O Lunges – 0-90°, Leg press – 0-90°
o Proprioception exercises
Begin use of the stationary bicycle Phase III (Weeks 13 16)
Phase III (Weeks 12-16)
Weight bearing: Full weight bearing with normal gait pattern Person of Matiers - Full (Painless ROM)
Range of Motion – Full/Painless ROM The second is Francisco
Therapeutic Exercises Continue with good and homestring strength aning.
Continue with quad and hamstring strengthening Fogus on single log strength
Focus on single-leg strengthBegin jogging/running
c , cc c.
O Plyometrics and sport-specific drills Phase IV (Months 4.6)
Phase IV (Months 4-6)
Gradual return to athletic activity as tolerated
Maintenance program for strength and endurance
Comments: Patients should avoid tibial rotation for 4-6 weeks post-op
Frequency: times per week Duration: weeks
Signature: Date: