

Professor of Orthopaedic Surgery Division of Sports Medicine Tel: (646) 501-7223

Proximal or Distal Realignment +/- Medial Patellofemoral Ligament Reconstruction Rehabilitation Program

Name:			Date:
Diagnosis:		-	Date of Surgery:
Pha	se I –In	nmediate Postoperative Phase (Day 1-5)	
— •	Goals:		
	0	Diminish swelling/inflammation (control hemar	throsis)
	0	Diminish postoperative pain	
	0	J 1 1	
	0	Independent ambulation	
•	Brace:		
	0	Brace for ambulation only (POD 1 to Week 4)	
•	_	t-bearing	
		As tolerated with two crutches (approx. 50% WE	3)
•		ng/Inflammation Control	
		Cryotherapy	
	0	Compression bandages	
	0	1 1	
•		of Motion	
		Full passive knee extension	
		Flexion to 45° (day 1-4)	
	0	Flexion to 60° (day 5)	
	0	3	
•	Flexibi	•	
	0	Hamstring and calf stretches	
	0	PROM/AAROM within ROM limitations	
Ph	ase II -	Acute Phase (Week 2-4)	
•	Goals:		
	0	Control swelling and pain	
	0	Promote healing of realignment tibial tuberosity	
	0	Quadriceps strengthening	
•	Brace		
	0	Continue brace for ambulation only	
	0	Discontinue brace (week 4)	
•	Weigh	t-bearing	
	0	Progress WBAT (2 crutches for 4 weeks)	
•	 Swelling/inflammation 		
	0	Continue use of cryotherapy	
	0	Compression bandage	
	0	Elevation	
•	Range	of motion	
	0	PROM/AAROM exercises	

ROM 0-75° (week 1-3)ROM 0-90° (week 4)

Michael J. Alaia, MD

Professor of Orthopaedic Surgery Division of Sports Medicine Tel: (646) 501-7223

- Muscle Retraining
 - o Electrical muscle stimulation to quads
 - Quad setting isometrics
 - Straight leg raises (flexion)
 - Hip adduction/abduction
 - Hip extension
 - o GENTLE submaximal isometric knee extension
 - o Week 4
 - Light leg press
 - Vertical squats (no weight)
- Flexibility
 - Continue hamstring, calf stretches

Phase III -Subacute Phase "Motion" Phase (Week 5-8)

- Goals
 - o Gradual improvement in ROM
 - Improve muscular strength and endurance
 - o Control forces on extension mechanism
- Weight-bearing
 - o One crutch (week 4-6)
 - o Discontinue crutch (week 6)
- Range of motion
 - o PROM 0-115°
 - o PROM 0-125°
 - o PROM 0-125/135°
- Exercises
 - o Continue electrical muscle stimulation to quadriceps
 - o Quadriceps setting isometric
 - o Hip adduction, abuction, and extension
 - Vertical squats
 - o Leg press
 - Knee extension light (0-60°)
 - o Bicycle (week 6-8)
 - o Pool program [walking, strengthening (when able)]
- Flexibility
 - Continue all stretching exercises for LE

Phase IV -Strengthening Phase (weeks 9-16)

- Criteria to Progress to Phase IV
 - o ROM at least 0-115 degrees
 - Absence of swelling/inflammation
 - Voluntary control of quads
- Goals
 - Gradual improvement of muscular strength
 - Functional activities/drills
- Exercises
 - o ½ vertical squats (0-60°)
 - Wall squats (0-60°)
 - Leg press
 - Forward lunges



Comments:

Frequency: ____ times per week

Signature:

Michael J. Alaia, MD

Date: _____

Professor of Orthopaedic Surgery Division of Sports Medicine Tel: (646) 501-7223

NYU LANG	Tel: (646) 501-7223	
0 0 0	Lateral lunges Lateral step-ups Knee extension (60-0°) Hip adduction/abduction Bicycle Stairmaster	
• Criteria	Return to Activity Phase a to enter Phase V Full non-painful ROM Appropriate strength level (80% or greater of contralateral leg) Satisfactory clinical exam Functional return to specific drills es Functional drills Strengthening exercises Flexibility exercises	

Duration: _____ weeks